



Information About Diet and IBD

The idea of modifying diet to influence inflammatory bowel disease (IBD) is something that has generated much interest from people with IBD, health care professionals and scientists over the past decade. This has led to dietary research that helps our understanding of the relationship between food and IBD and development of evidence-based dietary recommendations.

In general, most patients with IBD should follow the Australian Guide to Healthy Eating, and do not need to restrict their diet. However, there is emerging or established evidence to support specialty diets for specific situations. Outlined below are the dietary recommendations for: (1) when IBD is in remission, (2) when IBD is active, and (3) to treat complications of IBD. Determining whether you have active disease or remission is undertaken by endoscopy, imaging and/or blood and stool tests, but can also be indicated by symptoms. It is important to note that symptoms alone are not the best markers of active disease or remission. Many people with IBD will develop a gut sensitivity, meaning symptoms develop without active inflammation. These are 'functional' symptoms, like in irritable bowel syndrome (IBS). On the other hand, active disease can occur without causing many symptoms.

1. Diet when IBD is in remission

There is some preliminary evidence to suggest that dietary factors may assist in keeping remission for longer in people with IBD. Many of these factors are in line with the Australian Guide to Healthy Eating, which describes a well-balanced diet with minimal dietary restrictions beyond those recommended for general health and well-being. Including a good amount of fibre-containing foods is encouraged, unless reducing fibre is necessary due to strictures (narrowing of the bowel) being present (see section 3).

Does general dietary guidance differ for Crohn's disease and ulcerative colitis?

The broad dietary recommendations are the same for both Crohn's disease and ulcerative colitis. There are, however, some early data suggesting that eating more fibre and including fish a few times per week has a protective effect for Crohn's disease, whilst consuming less red and processed meats may have a protective effect in ulcerative colitis.

2. Diets to treat active IBD

During active disease, the goal of management is to heal the gut (induce remission) and then to keep the gut healed (maintain remission). Your treating doctor will guide you on the status of your disease and appropriate dietary treatments to both induce and maintain remission.

Crohn's disease

Exclusive Enteral Nutrition (EEN):

EEN is a nutritionally complete medical drink used to induce remission in Crohn's disease as an alternative to corticosteroids. This therapy involves solely consuming these drinks, usually for six to eight weeks. All food and fluids (except water) are removed from the diet. EEN is a treatment that needs to be commenced and monitored with a gastroenterologist and dietitian to ensure it is done correctly and safely.



Australian Guide to Healthy Eating


Enjoy a wide variety of nutritious foods from these five food groups every day.

Drink plenty of water.

Grain (cereal) foods, mostly wholegrain and/or high cereal fibre varieties



Vegetables and legumes/beans



Lean meats and poultry, fish, eggs, tofu, nuts and seeds and legumes/beans



Fruit



Milk, yoghurt, cheese and/or alternatives, mostly reduced fat



Use small amounts



Only sometimes and in small amounts



Crohn's Disease Exclusion Diet (CDED):

CDED is a diet that has been developed as an alternative to EEN and shown to be similar for inducing remission in children with mildly active Crohn's disease. This diet is intended for short term use and involves consuming a medical nutrition drink with an allowance of a limited number of prescribed foods. CDED needs supervision with a gastroenterologist and dietitian to ensure individual nutritional needs are met and the diet is having the intended treatment effect.

Ulcerative colitis

At this stage there is insufficient evidence to recommend a defined dietary strategy beyond healthy eating guidelines for treating active ulcerative colitis.

Complementary and alternative diets that claim to treat active IBD

There are many diets promoted on the internet, social media, and by some complementary and alternative health practitioners, claiming to treat IBD. Some of these diets include Specific Carbohydrate Diet, Paleo autoimmune protocol Diet, plant-based diet, and IBD anti-inflammatory diet.

Much of this information is conflicting, and many of these diets are not supported by sufficient scientific evidence to date. Though there may be some "success stories" that you may read or hear about, the experience of some individuals may not be safe or applicable for you.

Furthermore, improvement in symptoms may not reflect improvement in inflammation in the bowel. To accurately study the effect of diet on inflammation, studies should assess disease activity using endoscopy, imaging and/or blood and stool tests. The risks with some of these diets is that they could cause more harm and delay remission, lead to unnecessary dietary restriction and nutritional inadequacy. This topic can be confusing, so please discuss this with your IBD team.

3. Dietary strategies to treat common complications of IBD

Diet may be used to treat arising problems that commonly occur in people with IBD, as described below. ***It is recommended that everyone with IBD have an opportunity to meet with a dietitian to discuss and tailor their diet.*** An IBD specialist dietitian can guide specific dietary education and support for:

1. Treating malnutrition (under- and over-nutrition)

Both under- and over-nutrition (overweight or obesity) is common in people with IBD and can lead to fatigue, reduced quality of life, depression and may weaken response to medications. People with malnutrition are more likely to become sick and take longer to recover. This can also lead to increased risk of complications after surgery. Furthermore, impairment of nutrient absorption is commonly seen in active Crohn's disease and can lead to specific nutritional deficiencies. Beyond a healthy diet, there is limited evidence for nutritional supplementation in IBD, unless needed to correct specific nutritional deficiencies. Some people may need iron (orally or intravenous), vitamin B12 and/or vitamin D replacement. If you are concerned that you have malnutrition, it is recommended that you see a dietitian.

2. Managing symptoms of irritable bowel syndrome (IBS)

IBS describes symptoms that are from gut sensitivity and/or altered gut movement. This can be managed with dietary and non-dietary therapies such as a low or modified FODMAP diet, changing eating behaviour, or gut-directed hypnotherapy to help control symptoms.

3. Prevent bowel obstruction

Bowel obstruction (partial or complete) can occur in people with small intestinal Crohn's disease with a stricture (narrowing of the intestinal space where food and stool passes). If this is suspected or occurs, your gastroenterologist and dietitian can guide you on a modified fibre, low fibre or low residue diet and for how long it would be needed.

4. Nutritional optimisation before and/or after surgery

Having good nutritional status before undergoing surgery is important to encourage good surgical outcomes, such as a short hospital stay, fast recovery of bowel function and reducing risk of post-surgical complications. Often, medical nutrition drinks are recommended for 7-10 days leading up to planned surgery to optimise nutritional status to improve surgical outcomes. In people with Crohn's disease, a period of EEN (as described earlier) before planned surgery may also be recommended to both improve nutrition and reduce active disease. This may be used in preference to corticosteroids. After surgery, your surgeon will gradually allow you to restart oral intake. Some patients may have narrowing at the join of the bowel due to swelling, and require a short- term low fibre diet to allow the site to heal and bowel movements to start. Your surgeon and dietitian will guide the degree and timeframe for fibre restriction, if needed.

5. Diet for people with a stoma Most people with a stoma have an acceptable output and do not need to modify their diet. Generally, the Australian Guide to Healthy Eating is recommended. One of the roles of the large intestine is to reabsorb water, so ensuring good hydration by having plenty of fluid is important. You may benefit from further consultation with a specialist dietitian if your stoma output is too much, too watery, or too gassy. Please refer to the Life with a Stoma information sheet for further information.

6. Supplementary therapies – prebiotics and probiotics, curcumin, fish oil

There is little evidence to support the use of most commercial prebiotic (to stimulate growth of beneficial bacteria) or probiotic (live bacteria) supplements for most people with IBD. There may be specific situations where probiotics may be beneficial, including in people with an ileoanal pouch and recurrent pouchitis. Small studies suggest that curcumin may be helpful in people with ulcerative colitis but further studies are needed. Fish oil supplements are not likely to be beneficial for IBD but can be good for general health.

Acknowledgements

This resource was developed in 2021 by the GESA IBD Patient Information Materials Working Group that included the following health professionals: Mayur Garg (Chair, Gastroenterologist) Aysha Al-Ani (Gastroenterologist) George Alex (Gastroenterologist - Paediatric) Vinna An (Colorectal Surgeon) Jakob Begun (Gastroenterologist) Maryjane Bettehem (Stomal Therapy Nurse) Robert Bryant (Gastroenterologist) Britt Christensen (Gastroenterologist) Rosemary Clerehan (Educational Linguist) Susan Connor (Gastroenterologist) Sam Costello (Gastroenterologist) Basil D'Souza (Colorectal Surgeon) Alice Day (Senior Gastrointestinal Dietitian) Kevin Greene (Consumer Representative) Geoff Haar (IBD Pharmacist) Emma Halmos (Senior Gastrointestinal Dietitian) Tim Hanrahan (Gastroenterology Trainee) Heidi Harris (IBD Clinical Nurse Consultant) Katherine Healy (Senior Gastrointestinal Dietitian) Simon Knowles (Specialist Gastrointestinal Psychologist) Taryn Lores (Health Psychologist) Raphael Luber (Gastroenterologist) Antonina Mikocka-Walus (Specialist Gastrointestinal Psychologist) Marion O'Connor (IBD Clinical Nurse Consultant) Meera Rajendran (IBD Pharmacist) Clarissa Rentsch (IBD Pharmacist) Sally Stockbridge (CCA Consumer Representative) Julie Weldon (CCA Consumer Representative) Charys Winter (IBD Clinical Nurse Consultant)